



477 W. Horton Rd.  
 Bellingham, WA 98226  
 Phone (360) 933-4892  
 Fax (360) 933-1197

Patient Name: \_\_\_\_\_

Date of Birth: \_\_\_\_\_ Wt: \_\_\_\_\_ Ht: \_\_\_\_\_

IV Access: \_\_\_\_\_

Allergies: \_\_\_\_\_

### IV Immune Globulin (IVIG) Order Form

- ◆ Orders are initiated unless crossed out by provider.
- Check box to initiate order.

Please complete this form and fax to (360)933-1197

**Diagnosis:** \_\_\_\_\_ ICD-10: \_\_\_\_\_

**Prescreening:**

Has the patient previously received IVIG?  No  Yes, brand and date: \_\_\_\_\_  
 Past Medical History (RPh may recommend additional premedication):  
 Migraine  Thrombosis  Diabetes  Renal dysfunction

**Medication Orders:**

- ◆ IVIG (Brand will be selected by pharmacy, unless specified)  Specific brand (if medically necessary): \_\_\_\_\_
  - Dose: \_\_\_\_\_g/kg (final dose determined by Pharmacy) **OR** \_\_\_\_\_grams (total dose)
  - Frequency: Give IV every \_\_\_\_\_
  - Other instructions: \_\_\_\_\_

**PHARMACY USE**

Pharmacist will use actual body weight (ABW) or adjusted body weight (adjBW) if ABW is >30% more than the ideal body weight (IBW), unless prescriber indicates otherwise.

ABW = \_\_\_\_\_kg      IBW = \_\_\_\_\_kg      If applicable, adjBW (IBW+0.4[ABW-IBW]) = \_\_\_\_\_kg

**Final dosing weight:** \_\_\_\_\_

**Calculated IVIG dose: (round to nearest 5g or available vial size)** \_\_\_\_\_ RPh initial/date: \_\_\_\_\_

- ◆ Following manufacturer's recommendations, initiate infusion at low end of range. Increase slowly every 15 to 30 minutes if tolerated until entire dose is infused.
- ◆ Alteplase 2mg IV to declot central IV access per Infusion Solutions protocol as needed for occlusion.
- ◆ Flush line with D5W, 0.9% NaCl and/or Heparin 10 units/ml or 100 units/ml per Infusion Solutions protocol.
- ◆ Lidocaine 1% - up to 0.2ml intradermally PRN (may buffer with sodium bicarbonate 8.4% in 10:1 ratio).

**Premedication (15 to 30 minutes before infusion):**

- ◆ Diphenhydramine:  50mg IV  25mg IV
- ◆ Acetaminophen:  1000mg PO  500mg PO
- ◆ Other: \_\_\_\_\_

**To Manage Infusion Reactions:**

- Methylprednisolone 125mg IV x1 dose PRN severe urticaria, pruritis, or SOB
- ◆ Infusion Reaction Management per Infusion Solutions Protocol
- Other: \_\_\_\_\_

**Nursing Orders:**

- ◆ If no central IV access, RN to insert peripheral IV, rotate site every 72 to 120 hours or as needed.
- ◆ Obtain weight before each dose.
- ◆ Monitor vital signs (temp, HR, RR, BP) before therapy, every 15 min x1 hour, every hour, and at completion of infusion.
- ◆ If an infusion reaction occurs, decrease rate by 30 ml/hr every 15 minutes and monitor vital signs until symptoms subside. If reaction persists or worsens, stop infusion and notify physician.
- Other: \_\_\_\_\_

**Labs:**  Serum Creatinine (recommend at least every 6 months) every \_\_\_\_\_  
 \_\_\_\_\_ every \_\_\_\_\_

\_\_\_\_\_  
*Prescriber Signature*

\_\_\_\_\_  
*Date*

\_\_\_\_\_  
*Please Print Name*